

Primordial Fluctuations within Teleparallelism

Yi-Peng Wu^{1,3a} and Chao-Qiang Geng^{1,2,3b}

¹*Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300*

²*Physics Division, National Center for Theoretical Sciences, Hsinchu, Taiwan 300*

³*Kavli Institute for Theoretical Physics China (KITPC), CAS, Beijing 100190, China*

(Dated: January 4, 2012)

Abstract

We study the cosmological perturbations for the possible inflation scenario in the teleparallel description of general relativity. A 3+1 decomposition of the vierbein field, resulting in an identical formalism to the Hamiltonian formulation of TEGR under a certain time gauge, provides the primordial fluctuations of the single field inflationary model in the teleparallel formulation indistinguishable from the standard one. For the higher-order action, $f(T)$ gravity, we find that extra degrees of freedom can be fixed by the time gauge so that the same parametrization is applied. The equivalence between conformal frames may break down due to the Lorentz violating rescaling process.

^a E-mail address: s9822508@m98.nthu.edu.tw

^b E-mail address: geng@phys.nthu.edu.tw

I. INTRODUCTION

Inflation is one of the most successful cosmological models, which is able to explain the structure of the universe observed today, in particular, the flatness problem. The predictions of each inflationary model are obtained through the quantization of the primordial fluctuations. Such scenario is demonstrated in [1, 2] for the simplest case that gravity is minimally coupled to a scalar field. The discussion can be also applied to higher-order gravity actions which have conformal equivalences to the minimal coupling cases [1]. More generically, it can be used in the most general second-order field equations [3, 4] or general modified gravitational models of inflation [5, 6].

Recently, the gravity constructed within “distant parallelism” or “teleparallelism” has received much attention of cosmological interests for both the early universe [7] and the late time cosmic acceleration [8]. While the teleparallel equivalence of general relativity (TEGR) is, so far, indistinguishable from general relativity (GR), its high order generalization, $f(T)$ gravity, contains some novel features other than $f(R)$ gravity. In particular, the local Lorentz invariance cannot be preserved as some extra degrees of freedoms are found [9, 10]. This feature is explicit in the Einstein frame of $f(T)$ gravity theories as an exotic scalar-torsion coupling appears in the Lagrangian [11]. However, under homogeneous and isotropic principles in cosmology, the effects of the extra degrees of freedom need to proceed to the perturbed equations. It is shown in [12] that under a gauge-invariant analysis, one extra degree of freedoms causes severe constraints on $f(T)$ gravity models.

Here, we investigate the cosmological perturbations of some possible inflationary model under the teleparallel description of general relativity. A 3+1 decomposition is found for vierbein fields which can lead to the ADM formulation for the metric commonly used in the literature. After applying such decomposed vierbein to TEGR, we find that the resulting torsion scalar T is identical to the Hamiltonian formulation under a certain time gauge consideration [13]. The representation is convenient for comparison between the formulations of general relativity.

In this paper, we first demonstrate that TEGR shares the same quadratic actions as GR for both scalar and tensor perturbations. This implies that the single field inflationary models in the teleparallel formulation can satisfy current observations. We also discuss the application of the possible inflation scenario driven by the higher-order gravity in TEGR

similar to $f(R)$ theories. By imposing the certain time gauge condition, we find that some of extra degrees of freedom in the general $f(T)$ theories are fixed so that the background vierbein can be simply chosen. Although the cosmological perturbation of the time gauge constraint $f(T)$ theory can proceed under the same parametrization as TEGR, one should still address the subtleties on the Lorentz violating rescaling for the conformal transformation. For the *coordinate and tangent frames*, Greek indices $\mu, \nu, \dots$ and capital Latin indices $A, B, \dots$ run over space and time, while Latin indices $i, j, \dots$ and $a, b, \dots$, represent the spatial part of 1, 2, 3, respectively.

The paper is organized as follows. In Sec. II, we review teleparallel gravity. In Sec. III, we perform the quadratic computation in TEGR. We extend our discussion to the high order generalization of TEGR in Sec. IV. Finally, conclusions are given in Sec. V.

II. TELEPARALLEL GRAVITY

Originated from the Einstein's approach to unify gravitation with electromagnetism [14], teleparallel gravity takes the vierbein field $\mathbf{e}_A(x^\mu)$ as the dynamical variables, which are also orthonormal bases of the tangent space for each point x^μ on the manifold. These bases satisfy the relation $\mathbf{e}_A \cdot \mathbf{e}_B = \eta_{AB}$, where $\eta_{AB} = \text{diag}(1, -1, -1, -1)$, and are related to the coordinate bases ∂_μ via the components e_A^μ , *i.e.*, $\mathbf{e}_A = e_A^\mu \partial_\mu$. Therefore, the metric tensor is obtained from the dual vierbein as

$$g_{\mu\nu}(x) = \eta_{AB} e_A^\mu(x) e_B^\nu(x). \quad (2.1)$$

On the other hand, teleparallel gravity uses the curvatureless Weitzenböck connection, $\Gamma_{\mu\nu}^\rho = e_A^\rho \partial_\nu e_\mu^A$, to define its covariant derivative and the torsion tensor

$$T_{\mu\nu}^\rho = \Gamma_{\nu\mu}^\rho - \Gamma_{\mu\nu}^\rho = e_A^\rho (\partial_\mu e_\nu^A - \partial_\nu e_\mu^A). \quad (2.2)$$

The relation between the Weitzenböck connection and the torsionless Levi-Civita connection $\bar{\Gamma}_{\mu\nu}^\rho$ used in GR is given through the contorsion tensor $K_{\mu\nu}^\rho = \frac{1}{2}(T_{\nu\mu}^\rho + T_{\mu\nu}^\rho - T_{\mu\nu}^\rho)$ as

$$\Gamma_{\mu\nu}^\rho = \bar{\Gamma}_{\mu\nu}^\rho + K_{\mu\nu}^\rho. \quad (2.3)$$

As demonstrated in [15], the most general Lagrangian density, which is the quadratic of the torsion tensor, is of the form

$$\mathcal{L} = a_1 T_\rho^{\mu\nu} T_{\mu\nu}^\rho + a_2 T^{\mu\nu} T_{\mu\nu}^\rho + a_3 T_{\rho\mu}^\rho T_\nu^{\nu\mu} + a_0, \quad (2.4)$$

where a_i are free parameters. An equivalent description to GR within teleparallelism is established in [13] and can be simply taken as the Lagrangian (2.4) with the choice $a_1 = 1/8$, $a_2 = -1/4$ and $a_3 = -1/2$ [16]. Consequently, the Lagrangian density of TEGR is given by

$$\begin{aligned}\mathcal{L}_{TEGR} &= \frac{1}{2}T = \frac{1}{8}T_\rho^{\mu\nu}T_\mu^\rho - \frac{1}{4}T^{\mu\nu}_\rho T^\rho_{\mu\nu} - \frac{1}{2}T^\rho_{\rho\mu}T^\nu{}^\mu_\nu \\ &\equiv \frac{1}{2}S_\rho^{\mu\nu}T^\rho_{\mu\nu},\end{aligned}\quad (2.5)$$

with the definition $S_\rho^{\mu\nu} = \frac{1}{2}(K^{\mu\nu}_\rho + \delta_\rho^\mu T^{\alpha\nu}_\alpha - \delta_\rho^\nu T^{\alpha\mu}_\alpha)$. The field equation of TEGR is obtained by variation of the vierbein field e^μ_A with respect to the action $S = \frac{1}{2}\int d^4x eT + S_m$, given by

$$e^{-1}\partial_\mu(ee^\rho_A S_\rho^{\mu\nu}) - e^\lambda_A T^\rho_{\mu\lambda} S_\rho^{\nu\mu} - \frac{1}{4}e^\nu_A T = \frac{1}{2}e^\rho_A \Theta_\rho{}^\nu, \quad (2.6)$$

where $\Theta_\rho{}^\nu$ is the energy-momentum tensor of the matter. By using the relation (2.3), it is found that $G^\nu_A = 2e^{-1}\partial_\mu(ee^\rho_A S_\rho^{\mu\nu}) - 2e^\lambda_A T^\rho_{\mu\lambda} S_\rho^{\nu\mu} - \frac{1}{2}e^\nu_A T$ where $G_{\mu\nu} = e^\mu_A G^\nu_A$ is nothing but the Einstein tensor. Therefore, we can rewrite (2.6) in the covariant way as $G_{\mu\nu} = \Theta_{\mu\nu}$, which is completely the same geometrical formulation as the Einstein equation in GR.

III. QUADRATIC COMPUTATION

The computation of the second order action for the primordial fluctuations of the standard inflationary model

$$S = \frac{1}{2}\int d^4x \sqrt{-g}[R + (\nabla\phi)^2 - 2V(\phi)] \quad (3.1)$$

has been reviewed nicely in [1] as well as [2] with a scenario commonly used in modified gravity theories which we shall follow in the rest of our discussion.

In order to make a clear comparison to the standard results, we consider the existence of a scalar field with a general potential in the Lagrangian of TEGR, given by

$$S = \frac{1}{2}\int d^4x e[T + (\nabla\phi)^2 - 2V(\phi)], \quad (3.2)$$

where $e \equiv \sqrt{-g}$ and the torsion scalar is defined as [16]

$$\begin{aligned}T &= \frac{1}{4}T_\rho^{\mu\nu}T^\rho_{\mu\nu} - \frac{1}{2}T^{\mu\nu}_\rho T^\rho_{\mu\nu} - T^\rho_{\rho\mu}T^\nu{}^\mu_\nu \\ &= R + 2\bar{\nabla}^\nu T^\mu_{\mu\nu}.\end{aligned}\quad (3.3)$$

The second equation of (3.3) is obtained from the mathematical manipulation by using the relation between Weitzenböck and Levi-Civita connections showing that T and R only differ by a divergent term with respect to the Levi-Civita derivative [9].

When we choose the vierbein to be $e_\mu^A = \text{diag}(1, a, a, a)$, (3.2) shares the some background equations with (3.1) in the flat FRW background:

$$3H^2 = \frac{1}{2}\dot{\phi}^2 + V, \quad \dot{H} = -\frac{1}{2}\dot{\phi}^2, \quad \text{and} \quad 0 = \ddot{\phi} + 3H\dot{\phi} + V_{,\phi}. \quad (3.4)$$

Notice that such a simple vierbein choice is an exact solution of theTEGR field equation corresponding to the flat FRW Einstein equation followed by the discussion in [15], and one can refer to [17] for the cases of open and closed universe.

A. ADM Decomposition

The computation of the quadratic action is convenient to proceed in the ADM formalism. As has been demonstrated in [12], we make the 3+1 decomposition of the vierbein with respect to a unit normal vector $e_\mu^0 = u_\mu$ in the frame such that $u_\mu = (N, \mathbf{0})$. The vierbein is found as:

$$\begin{aligned} e_\mu^0 &= (N, \mathbf{0}) \quad , \quad e_\mu^a = (N^a, h_a^i) \\ e_0^\mu &= (1/N, -N^i/N) \quad , \quad e_a^\mu = (0, h_a^i), \end{aligned} \quad (3.5)$$

where $N^i \equiv h_a^i N^a$ with $h_a^i h_b^i = \delta_b^a$ for the induced 3-vierbein h_a^i .

It is straightforward to define the 3-covariant derivative D_i with respect to the 3-Weitzenböck connections ${}^{(3)}\Gamma_{jk}^i = h_a^i \partial_k h_j^a$, which can satisfy the relation

$$D_i h_{jk} = 0 \quad ; \quad D_i N^j = h_a^j \partial_i N^a, \quad (3.6)$$

where h_{ij} is the induced 3-metric corresponding to the metric obtained from (3.5):

$$ds^2 = N^2 dt^2 - h_{ij}(dx^i + N^i dt)(dx^j + N^j dt). \quad (3.7)$$

The non-vanished torsions in this representation are

$$\begin{aligned} T_{j0}^0 &= \partial_j N/N, \\ T_{j0}^i &= D_j N^i - \frac{N^i}{N} \partial_j N - h_a^i \partial_0 h_j^a, \\ T_{jk}^i &= h_a^i (\partial_j h_k^a - \partial_k h_j^a) \equiv {}^{(3)}T_{jk}^i, \end{aligned} \quad (3.8)$$

where the last equation is referred to as the definition of the induced 3-torsion.

It is also found to be convenient to define the “extrinsic torsion” as

$$\Sigma_{ij} = \frac{1}{2N}(\dot{h}_{ij} - D_i N_j - D_j N_i) \quad (3.9)$$

in the following calculations. Note that the extrinsic curvature is given by $\bar{\Sigma}_{ij} = \frac{1}{2N}(\dot{h}_{ij} - \bar{D}_i N_j - \bar{D}_j N_i)$ with $\bar{D}_i$ the 3-Levi-Civita covariant derivative. We can derive the relation between extrinsic torsion and extrinsic curvature if further applying the relation (2.3) among the induced 3-connections:

$${}^{(3)}\Gamma^i_{jk} = {}^{(3)}\bar{\Gamma}^i_{jk} + {}^{(3)}K^i_{jk}, \quad (3.10)$$

so that

$$\Sigma_{ij} = \bar{\Sigma}_{ij} - \frac{N^k}{2N}(T_{ijk} + T_{jik}). \quad (3.11)$$

Under the decomposition (3.5), the torsion scalar (3.3) is obtained in terms of the extrinsic torsion and 3-torsion as

$$T = \Sigma_{ij}\Sigma^{ij} - (\Sigma + \frac{N^k}{N}T^i_{ik})^2 - T^{(3)} - 2\frac{\partial_k N}{N}T^i_{ik} + 2\frac{N^k}{N}T_{ijk}\Sigma^{ij} + \frac{N^k N^l}{2N^2}T^{ij}_k(T_{ijl} + T_{jil}), \quad (3.12)$$

where we define that $T^{(3)} = \frac{1}{4}T_l^{mn}T^l_{mn} - \frac{1}{2}T^{mn}_l T^l_{mn} - T^j_{jk}T^l_{lk}$. In order to further compare with the usual ADM formalism obtained in GR, we insert the relation (3.11) into (3.12). Using the differential by part $\bar{D}_k(NT^i_{ik}) = N\bar{D}_k T^i_{ik} + T^i_{ik}\partial_k N$ for the substitution, we have the final expression, given by

$$T = \bar{\Sigma}_{ij}\bar{\Sigma}^{ij} - \bar{\Sigma}^2 + R^{(3)} + \mathcal{D}_T, \quad (3.13)$$

with $\mathcal{D}_T = \frac{-2}{N}\bar{D}_k(NT^i_{ik})$ becoming a total divergence term in the action. Here, $R^{(3)}$ is obtained from the same definition as (3.3) with the relation $T^{(3)} = R^{(3)} + 2\bar{D}_j T^{lj}_l$.

Replacing $\bar{\Sigma}^{ij}$ by $\Pi^{ij} = \sqrt{h}(\bar{\Sigma}^{ij} - \bar{\Sigma}h^{ij})$ as the conjugate momentum of h_{ij} [19], we find that (3.13) becomes identical to Eq. (10) in [20], which is the 3+1 formulation of TEGR under the Schwinger’s time gauge consideration [21]. As a result, the condition $e^0_\mu = u_\mu$ for the decomposition (3.5) can be realized as a specific choice, required to lock the time axes of tangent frames to the general coordinate time axis. Note that such time gauge is not a necessary condition for the general 3+1 formulation of TEGR, which is different from the structure of the standard ADM formalism [22].

It is easy now to make a comparison with the ADM decomposition of Ricci scalar R [1]:

$$R = \bar{\Sigma}_{ij}\bar{\Sigma}^{ij} - \bar{\Sigma}^2 + R^{(3)} + \mathcal{D}_R, \quad (3.14)$$

where $\mathcal{D}_R = \frac{2}{N\sqrt{h}}\partial_t(\sqrt{h}\bar{\Sigma}) - \frac{2}{N}\bar{D}_i(\bar{\Sigma}N^i + h^{ij}\partial_j N)$. We find that the difference between T and R is given by the total divergence $\mathcal{D}_T - \mathcal{D}_R = 2\bar{\nabla}^\nu T^\mu_{\mu\nu}$.

B. Scalar Perturbations

We can now rewrite the action (3.2) via (3.13) as

$$S = \frac{1}{2} \int \sqrt{h} \left[N(\bar{\Sigma}_{ij}\bar{\Sigma}^{ij} - \bar{\Sigma}^2) + NR^{(3)} + N^{-1}(\dot{\phi} - N^i\partial_i\phi)^2 - Nh^{ij}\partial_i\phi\partial_j\phi - 2NV \right] + \mathcal{D}_T, \quad (3.15)$$

where h^a_i and ϕ play the role of dynamical variables in which one can choose a gauge to fix the time and spatial reparametrizations. We denote the first order quantities, ζ and γ , to parametrize the scalar and tensor perturbations, and use the gauge convenient for studying the quadric action as

$$\delta\phi = 0, \quad h^a_i = ae^\zeta(\delta^a_i + \frac{1}{2}\gamma^a_i), \quad (3.16)$$

where γ^a_i in general contains symmetric and antisymmetric parts, *i.e.*, $\gamma^a_i = \mathbf{s}^a_i + \mathbf{a}^a_i$. The antisymmetric part $\mathbf{a}^a_i$ is a distinct feature for tensor perturbations in teleparallelism. However, it does not contribute to the quadric calculation in the discussion given later. We define that $\gamma_{ij} = \eta_{ab}(\delta^a_i\gamma^b_j + \delta^b_j\gamma^a_i)/2$ with $\eta_{ab} = \text{diag}(-1, -1, -1)$ and $\gamma_{ii} = \partial_i\gamma_{ij} = 0$, leading to the induced metric to be

$$h_{ij} = a^2 e^{2\zeta}(\delta_{ij} + \gamma_{ij} + \frac{1}{4}\gamma_{ai}\gamma^a_j). \quad (3.17)$$

Temporally, we focus on the scalar quantity ζ and take $\gamma = 0$ in (3.17) to simplify the calculations. With in mind that N and N^i (or N^a) are as Lagrange multipliers, we perform the variations δN and $\delta N^i = h^i_a \delta N^a$ to (3.15), and obtain the constraints

$$\begin{aligned} h^i_a \bar{\nabla}^j [\bar{\Sigma}_{ij} - h_{ij}\bar{\Sigma}] &= 0, \\ R^{(3)} - (\bar{\Sigma}_{ij}\bar{\Sigma}^{ij} - \bar{\Sigma}^2) - N^{-2}\dot{\phi}^2 - 2V &= 0. \end{aligned} \quad (3.18)$$

By setting that $N^i = \partial_i\psi + N^i_T$ with $\partial_i N^i_T = 0$ and $N = 1 + N_1$ as suggested in [2], we derive the solutions

$$N_1 = \frac{\dot{\zeta}}{H}, \quad N^i_T = 0, \quad \psi = -\frac{\zeta}{a^2 H} + \chi, \quad \partial^2\chi = \frac{\dot{\phi}^2}{2H^2}\dot{\zeta}. \quad (3.19)$$

After substituting (3.19) back to (3.15), the Lagrangian is expressed in terms of ζ to the

second order

$$S_\zeta = \frac{1}{2} \int dt dx^3 \left\{ a e^\zeta \left(1 + \frac{\dot{\zeta}}{H} \right) [4\partial^2 \zeta - 2(\partial \zeta)^2 - 2a^2 V e^{2\zeta}] \right. \\ \left. + a^3 e^{3\zeta} \frac{1}{1 + \frac{\dot{\zeta}}{H}} [-6(H + \dot{\zeta})^2 + \dot{\phi}^2] \right\}, \quad (3.20)$$

which is identical to the expression by using the background equations (3.4):

$$S_\zeta = \frac{1}{2} \int dt d^3 x a^3 \frac{\dot{\phi}^2}{H^2} (\dot{\zeta}^2 - a^{-2}(\partial \zeta)^2). \quad (3.21)$$

This final expression for S_ζ is identical to the result of the theory (3.1).

C. Tensor Perturbations

As demonstrated in [2], the γ^2 term and the scalar-tensor crossing terms appear only in the cubic calculations of the Lagrangian, which are beyond the main concern of this paper. Hence, we simply set $\zeta = 0$ in (3.17) and consider the contribution only from γ_i^a . After inserting to (3.15) the quadratic action is obtained as

$$S_\gamma = \frac{1}{8} \int dt d^3 x a^3 [(\dot{\gamma}_{ij})^2 - a^{-2}(\partial_i \gamma_{jk})^2]. \quad (3.22)$$

Note that, despite the tensor fluctuation γ_i^a contains some extra components, i.e., the anti-symmetric part $\mathbf{a}_i^a$ comparing to GR. The contribution of $\mathbf{a}_i^a$ does not show up at this level.

We can define $\delta_{ia} \mathbf{s}_j^a = \mathbf{s}_{ij}$ and $\delta_{ia} \mathbf{a}_j^a = \mathbf{a}_{ij}$, so that actually

$$\gamma_{ij} = \frac{1}{2} \eta_{ab} (\delta_i^a \gamma_j^b + \delta_j^b \gamma_i^a) = \mathbf{s}_{ij}. \quad (3.23)$$

From the fact that (3.21) and (3.22) are the same results as the theory (3.1), we simply conclude that the changing of the dynamical variable from h_{ij} to h_i^a unchanges the propagating degrees of freedom in linear perturbations as the additional continent $\mathbf{a}_i^a$ drops out in quadratic computations. Therefore, we have two tensor modes plus one scalar mode in teleparallel gravity, and the effects from $\mathbf{a}_i^a$ in higher order perturbations worth for further investigations.

IV. HIGH ORDER GENERALIZATION

The high order generalization of TEGR was first proposed in [7] as an alternative inflation model. As analogy to $f(R)$ gravity, the study on the modification of the torsion scalar T

can be generalized to an arbitrary function with the action

$$S = \frac{1}{2} \int dx^4 e f(T). \quad (4.1)$$

In order to study the quadratic action of the high order Lagrangian, it is provided to be convenient to perform the conformal transformation [1]. We simply review the transformation of $f(T)$ gravity to the Einstein frame first and one can find the detail in [11]. We use the formulation

$$f(T) = FT - 2V, \quad F \equiv \frac{df}{dT}, \quad V = \frac{FT - f(T)}{2}, \quad (4.2)$$

and define $\hat{e}_\mu^A = \sqrt{F} e_\mu^A \equiv \Omega e_\mu^A$. The torsion scalar is transformed as

$$T = \Omega^2 [\hat{T} - 4\hat{\partial}^\mu \omega \hat{T}^\rho_{\rho\mu} + 6(\hat{\nabla} \omega)^2], \quad (4.3)$$

where $\hat{\partial}^\mu \omega \equiv \hat{\partial}^\mu \Omega / \Omega$. Since $e = \Omega^{-4} \hat{e}$ and $F = \Omega^2$, the action (4.1) is rewritten as

$$S = \frac{1}{2} \int dx^4 \hat{e} \left[\hat{T} - \frac{4}{\sqrt{6}} \hat{\partial}^\mu \varphi \hat{T}^\rho_{\rho\mu} + (\hat{\nabla} \varphi)^2 - 2U(\varphi) \right], \quad (4.4)$$

where we have defined that $d\varphi = \sqrt{6} d\omega = \sqrt{6} dF / 2F$ and $U = V / F^2$. Comparing the action (4.4) with (3.2), there is an additional scalar-torsion coupling term, $\hat{\partial}^\mu \varphi \hat{T}^\rho_{\rho\mu}$, which breaks the local Lorentz invariance and gives the extra degrees of freedom. The corresponding field equation of (4.4) is obtained by variation with respect to the vierbein field $\hat{e}_\mu^A$, where the covariant representation is

$$G_{\mu\nu} = \Theta_{\mu\nu}^\varphi + H_{\mu\nu} \quad (4.5)$$

with $\Theta_{\mu\nu}^\varphi$ the corresponding energy-momentum tensor of the scalar field φ , and $H_{\mu\nu}$ is obtained from variation of the coupling $\hat{e} \hat{\partial}^\mu \varphi \hat{T}^\rho_{\rho\mu}$, given by:

$$\begin{aligned} \Theta_{\mu\nu}^\varphi &= \hat{\partial}_\mu \varphi \hat{\partial}_\nu \varphi - \frac{1}{2} g_{\mu\nu} (\hat{\nabla} \varphi)^2 + g_{\mu\nu} U \\ H_{\mu\nu} &= \frac{2}{\sqrt{6}} [\hat{g}_{\mu\nu} \hat{\partial}^\lambda \varphi \hat{T}^\rho_{\rho\lambda} - \hat{\partial}^\lambda \varphi \hat{T}_{\nu\mu\lambda} - 2\hat{\partial}_\nu \varphi \hat{T}^\rho_{\rho\mu}] - \frac{2}{\sqrt{6}} \hat{e}^{-1} \hat{g}_{\nu\lambda} \hat{e}_\mu^A \hat{\partial}_\alpha [\hat{e} (\hat{\partial}^\lambda \varphi e_A^\alpha - \hat{\partial}^\alpha \varphi e_A^\lambda)] \end{aligned} \quad (4.6)$$

respectively.

It is easy to observe that $H_{\mu\nu}$ in general is not a symmetric tensor while $G_{\mu\nu}$ and $\Theta_{\mu\nu}^\varphi$ are. Therefore, following the discussion in [9] but here in the Einstein frame, the field equation (4.5) can be splitted into:

$$G_{\mu\nu} = \Theta_{\mu\nu}^\varphi + H_{(\mu\nu)} \quad (4.7)$$

$$0 = H_{[\mu\nu]} \quad (4.8)$$

where $H_{\mu\nu} = H_{(\mu\nu)} + H_{[\mu\nu]}$. The presence of (4.8) makes (4.5) an equation for all 16 components instead of 10, indicating that the theory (4.4) in fact precesses more degrees of freedoms than (3.1). We can find from (4.8) that extra degrees of freedoms are fully caused by the scalar-torsion coupling $\hat{\partial}^\mu \varphi \hat{T}^\rho_{\rho\mu}$. As a result, the specific background choice $\hat{e}_\mu^A = \text{diag}(1, \hat{a}, \hat{a}, \hat{a})$ with $\hat{a} = \Omega a$ can not be used directly for parametrizing the linear perturbation as the minimal coupled TEGR.

A. $f(T)$ Theory with Schwinger's Time Gauge

In the non-linear formulation of teleparallel gravity, the lack of local Lorentz invariance introduces extra degrees of freedom so that the time axes of tangent frames can not be locked exactly to the time axis of the general coordinate [12]. In order to describe the flat FRW background by the simple choice $\hat{e}_\mu^A = \text{diag}(1, \hat{a}, \hat{a}, \hat{a})$ we impose the Schwinger's time gauge condition [21], which will reduce the degrees of freedom in the generic $f(T)$ theories.

To see this, we decompose the vierbein field $\hat{e}_\mu^A$ similar to (3.5) to obtain the quadratic action of (4.4). The torsion scalar $\hat{T}$ is found in (3.13) where we denote $\hat{\Sigma}_{ij}$ as the extrinsic curvature with respect to the induced metric $\hat{h}_{ij}$. For simplicity, we represent the scalar-torsion coupling under the integration by part as

$$\hat{e}(\varphi \hat{\nabla}^\mu \hat{T}^\rho_{\rho\mu}) = -\varphi \partial_t(\sqrt{\hat{h}} \hat{\Sigma}) + \sqrt{\hat{h}} \varphi \bar{D}_i(N^i \hat{\Sigma} + h^{ij} \partial_j N - N \hat{T}^j_i), \quad (4.9)$$

where the decomposed $\hat{\nabla}^\mu \hat{T}^\rho_{\rho\mu}$ can be found in the previous discussion. Even without knowing the background field, it is explicit that the term $\sqrt{\hat{h}} \varphi \bar{D}_i(\dots)$ will become a total divergence if we also impose the condition $\delta\varphi = 0$ where φ preserves only its background value $\varphi = \varphi(t)$. The action (4.4) under this condition is written as

$$S = \frac{1}{2} \int \sqrt{\hat{h}} \left[N(\hat{\Sigma}_{ij} \hat{\Sigma}^{ij} - \hat{\Sigma}^2) + N \hat{R}^{(3)} + \frac{4}{\sqrt{6}} \dot{\varphi} \hat{\Sigma} + N^{-1} \dot{\varphi}^2 - 2NU \right], \quad (4.10)$$

where we have neglected two total divergence terms, $\mathcal{D}_{\hat{T}}$ from the decomposed $\hat{T}$ and $\sqrt{\hat{h}} \varphi \bar{D}_i(\dots)$ as mentioned in (4.9).

One may observe that all terms inside (4.10) are no more than variables of the metric $\hat{h}_{ij}$ with its corresponding Levi-Civita derivative $\hat{D}_i$, and without higher than second-order time derivative. The change of variables from $\hat{h}_{ij}$ to $\hat{h}_i^a$ shows no difference in quadratic

computations for $\hat{\Sigma}_{ij}$ and $\hat{R}^{(3)}$ according to the discussion in Sec. III, while the only non-metric quantity $\hat{T}_j^{ji}$ drops as the total divergence. Hence, the degrees of freedom of (4.10) is the same as the action (3.15), which is found to have one scalar mode and two tensor modes in linear perturbations.

Remarkably, the non-metric 3-torsion tensor $\hat{T}_j^{ji}$ is also the only Lorentz violating term concerning the local Lorentz transformation $\hat{e}_{\nu}^{\tilde{A}} = \Lambda_{\tilde{B}}^{\tilde{A}}(x)\hat{e}_{\nu}^{\tilde{B}}$ to the Einstein frame vierbein field $\hat{e}_{\nu}^A$. It is easy to see from the fact that the metric is invariant under local Lorentz transformation since $\eta_{\tilde{C}\tilde{D}} = \Lambda_{\tilde{C}}^A \Lambda_{\tilde{D}}^B \eta_{AB}$ [16]. In general, T is not a local Lorentz scalar in the aspect of the original frame e_{μ}^A so that $f(T)$ does not preserve Lorentz invariance [9]. It is found in [10] that $f(T)$ gravity contains three extra degrees of freedom than GR from the analysis in the original Jordan frame (4.1). However, after applying the Schwinger's time gauge condition we find that the action (4.10) becomes, naively, similar to the structure of GR with some non-minimal derivative coupling to the scalar field [18]. The background $\hat{e}_{\mu}^A = \text{diag}(1, \hat{a}, \hat{a}, \hat{a})$ thereby can be more properly used to simplify the discussion on $f(T)$ theories under the constraint of the time gauge.

B. Quadratic Computations: A general overview

We can obtain the Hamiltonian and momentum constraints of (4.10) from the variation of N and N^a , written as

$$h_a^j \hat{D}^i \left[\hat{\Sigma}_{ij} - h_{ij} \hat{\Sigma} + h_{ij} \frac{4\dot{\varphi}}{\sqrt{6}N} \right] = 0, \quad (4.11)$$

$$\hat{R}^{(3)} - (\hat{\Sigma}_{ij} \hat{\Sigma}^{ij} - \hat{\Sigma}^2) - N^{-1} \frac{4}{\sqrt{6}} \dot{\varphi} \hat{\Sigma} - N^{-2} \dot{\varphi}^2 - 2U = 0, \quad (4.12)$$

respectively. Following the background $\hat{e}_{\mu}^A = \text{diag}(1, \hat{a}, \hat{a}, \hat{a})$, we can use the same parametrization as (3.16), and the field equations (4.5) and equation motion of φ are

$$\begin{aligned} 3H^2 &= \frac{1}{2} \dot{\varphi}^2 + U + \sqrt{6} \dot{\varphi} H \\ \dot{H} &= -\frac{1}{2} \dot{\varphi}^2 + \frac{1}{\sqrt{6}} \ddot{\varphi} - \frac{\sqrt{6}}{2} H \dot{\varphi} \\ 0 &= \ddot{\varphi} + 3H \dot{\varphi} + U' - \sqrt{6} H \frac{\dot{\varphi}}{\varphi}. \end{aligned} \quad (4.13)$$

We denote the perturbations as $N = 1 + N_1$ and $N^i = \partial_i \psi + N_T^i$, and hide the notation of the conformal frame in convenience for the following discussion. The quadratic action of

(4.10) is given by

$$\begin{aligned}
S_\zeta = & \frac{1}{2} \int dt dx^3 \left\{ a e^\zeta (1 + N_1) \left[-4\partial^2 \zeta - 2(\partial \zeta)^2 - 2a^2 U e^{2\zeta} \right] \right. \\
& + a^3 e^{3\zeta} \frac{1}{1 + N_1} \left[-6(H + \dot{\zeta})^2 + \dot{\phi}^2 + 4(H + \dot{\zeta})\partial^2 \psi \right] \\
& \left. + a^3 e^{3\zeta} \frac{\dot{\phi}}{1 + N_1} \frac{4}{\sqrt{6}} \left[3(H + \dot{\zeta}) - \partial^2 \psi \right] \right\}. \tag{4.14}
\end{aligned}$$

After some simple manipulation together with the constraint (4.11) and background equations (4.13), we find (4.14) can be rewritten as

$$\begin{aligned}
S_\zeta = & \frac{1}{2} \int dt dx^3 a^3 \left[-6G_T \dot{\zeta}^2 - 2F_T (\partial \zeta)^2 + 2\Xi N_1^2 \right. \\
& \left. - 4\Theta N_1 \partial^2 \psi + 4G_T \dot{\zeta} \partial^2 \psi + 12\Theta N_1 \dot{\zeta} - 4G_T N_1 \partial^2 \psi \right], \tag{4.15}
\end{aligned}$$

where $G_T = F_T = 1$, $\Xi = -U$ and $\Theta = H(1 - \dot{\phi}/\sqrt{6}H)$. The action (4.15) is in fact the representative of the generic quadratic action with the second-order field equations given in [3]. Following the discussion therein, we can arrive directly at the final expressions for tensor and scalar perturbations as

$$S_\gamma = \frac{1}{8} \int dt d^3x a^3 \left[G_T (\dot{\gamma}_{ij})^2 - a^{-2} F_T (\partial_i \gamma_{jk})^2 \right], \tag{4.16}$$

$$S_\zeta = \frac{1}{2} \int dt d^3x a^3 \left[G_S \dot{\zeta}^2 - a^{-2} G_S (\partial \zeta)^2 \right], \tag{4.17}$$

respectively, where

$$F_S = \frac{1}{a} \frac{d}{dt} \left(\frac{a}{\Theta} G_T^2 \right) - F_T, \tag{4.18}$$

$$G_S = \frac{\Xi}{\Theta^2} G_T^2 + 3G_T. \tag{4.19}$$

We find that tensor perturbations are unchanged from (3.22) despite the presence of the scalar-torsion coupling in (4.10), and scalar perturbations for any certain $f(T)$ inflation model shall satisfy the conditions of $F_S > 0$ and $G_S > 0$ to avoid ghost and instabilities [3].

C. Lorentz Violation Revisit

Although $f(T)$ theories can be converted as a single field non-minimally coupled to gravity through the time gauge, we wish to point out that (4.10) may still suffer by Lorentz violating subtlety despite its invariant under the transformation $\hat{e}_\nu^{\tilde{A}} = \Lambda_{\tilde{B}}^{\tilde{A}}(x) \hat{e}_\nu^{\tilde{B}}$. Implicitly, Lorentz violation is hidden through the conformal transformation as the conformal factor $\Omega = \sqrt{F}$

being a Lorentz-violating scalar to the original frame e_μ^A . From this point of view we find that $\hat{R}$, φ and U are all Ω involved and are not local Lorentz scalar either. One may speculate the possible consequence for the Lorentz violating scalar field φ , and the same question can also be found in the scalar-tensor formulations of $f(T)$ theories [23].

The issue on the equivalence between Jordan frame and Einstein frame becomes crucial for $f(T)$ theory due to the Lorentz violating rescaling process, $\hat{e}_\mu^A = \Omega e_\mu^A$. The “equivalence” is provided by including the running unit properly in the conformal frame [24]. Hence, it becomes considerable if given any unit redefinition needed for $f(T)$ theory from the Lorentz-violating conformal factor Ω .

V. CONCLUSIONS

In order to study the primordial behavior for gravity constructed under teleparallelism, especially for TEGR, we have performed a 3+1 decomposition of the vierbein field. The corresponding metric obtained from such decomposed vierbein is the ADM formulation commonly used in the generic models of inflation. We have found that the torsion scalar T under this representation differs from the decomposed Ricci scalar R by a total divergence, which is consistent with the general relation between T and R . Meanwhile, this decomposition can also be interpreted as a specific Hamiltonian formulation under the Schwinger’s time gauge condition. The quadratic computation of T can be taken as a rephrase of the standard result by changing the variable from h_{ij} to h_i^a . The reason is that linear perturbations give identical formulations to metric variables while all non-metric terms are dropped as the total divergence. Although in general the tensor part of vierbein contains extra components, its effects do not show up at the quadratic level, and the quantization process afterward will be no difference from the previous works.

The high order generalization of TEGR may also be considered as an alternative scenario for inflation by analogy to $f(R)$ theories. To discuss the cosmological perturbations, one needs to address first extra degrees of freedom introduced by the Lorentz violation in non-linear formulation of teleparallel gravity. Although we have shown that, in the conformal frame of $f(T)$ theory, the time gauge can fix the Lorentz issue and degrees of freedom simultaneously, the problem may convert to the Lorentz violating rescaling process which will break the equivalence between conformal frames, even at classical level.

ACKNOWLEDGMENTS

We are grateful to E. N. Saridakis and T. Kobayashi for the inspiring and helpful discussions. We thank KITPC for the wonderful programs of “Dark Matter and New Physics” and “String Phenomenology and Cosmology”. This work was partially supported by National Center of Theoretical Science and National Science Council (NSC-98-2112-M-007-008-MY3) of R.O.C.

-
- [1] V. F. Mukhanov, H. A. Feldman and R. H. Brandenberger, Phys. Rep. 215, 203 (1992).
 - [2] J. M. Maldacena, JHEP 0305, 013 (2003).
 - [3] T. Kobayashi, M. Yamaguchi, J. Yokoyama, arXiv:1105.5723 [hep-th].
 - [4] T. Kobayashi, M. Yamaguchi, J. Yokoyama, Phys. Rev. Lett. 105, 231302 (2010); X. Gao, T. Kobayashi, M. Yamaguchi, J. Yokoyama, arXiv:1108.3513 [astro-ph.CO]; A. D. Felice, S. Tsujikawa, Phys. Rev. D84, 083504 (2011).
 - [5] A. D. Felice, S. Tsujikawa, JACP 04, 029 (2011).
 - [6] X. Gao, D. A. Steer, [arXiv:1107.2642 [astro-ph.CO]]; X. Gao, JCAP **1110**, 021 (2011) [arXiv:1106.0292 [astro-ph.CO]].
 - [7] R. Ferraro and F. Fiorini, Phys. Rev. D75, 084031 (2007); Phys. Rev. D78, 124019 (2008).
 - [8] G. R. Bengochea and R. Ferraro, Phys. Rev. D **79**, 124019 (2009); E. V. Linder, Phys. Rev. D **81**, 127301 (2010); R. Myrzakulov, arXiv:1006.1120 [gr-qc]; P. Wu, H. W. Yu, Eur. Phys. J. **C71**, 1552 (2011); K. Bamba, C. Q. Geng, C. C. Lee, arXiv:1008.4036 [astro-ph.CO]; R. Zheng, Q. G. Huang, JCAP **1103**, 002 (2011); K. Bamba, C. Q. Geng, C. C. Lee, L. W. Luo, JCAP **1101**, 021 (2011); T. Wang, Phys. Rev. **D84**, 024042 (2011); K. K. Yerzhanov, S. R. Myrzakul, I. I. Kulnazarov and R. Myrzakulov, arXiv:1006.3879 [gr-qc]; P. Wu, H. W. Yu, Phys. Lett. **B693**, 415 (2010); G. R. Bengochea, Phys. Lett. **B695**, 405 (2011); P. Wu, H. W. Yu, Phys. Lett. **B692**, 176 (2010); S. H. Chen, J. B. Dent, S. Dutta and E. N. Saridakis, Phys. Rev. D **83**, 023508 (2011); J. B. Dent, S. Dutta, E. N. Saridakis, JCAP **1101**, 009 (2011); X. c. Ao, X. z. Li, P. Xi, Phys. Lett. **B694**, 186 (2010) arXiv:1010.4117 [gr-qc]; Y. Zhang, H. Li, Y. Gong, Z. H. Zhu, JCAP **1107**, 015 (2011); Y. F. Cai, S. H. Chen, J. B. Dent, S. Dutta, E. N. Saridakis, arXiv:1104.4349 [astro-ph.CO]. R. Ferraro, F. Fiorini, Phys. Lett. **B702**, 75 (2011);

- S. Chattopadhyay, U. Debnath, Int. J. Mod. Phys. **D20**, 1135 (2011); M. Sharif, S. Rani, Mod. Phys. Lett. **A26**, 1657 (2011); H. Wei, X. P. Ma, H. Y. Qi, Phys. Lett. **B703**, 74 (2011); R. Ferraro, F. Fiorini, arXiv:1106.6349 [gr-qc]; R. X. Miao, M. Li, Y. G. Miao, arXiv:1107.0515 [hep-th]; C. G. Boehmer, A. Mussa, N. Tamanini, arXiv:1107.4455 [gr-qc]; H. Wei, H. Y. Qi, X. P. Ma, arXiv:1108.0859 [gr-qc]; S. Capozziello, V. F. Cardone, H. Farajollahi, A. Ravanpak, arXiv:1108.2789 [astro-ph.CO]; P. Wu, H. Yu, Phys. Lett. **B703**, 223 (2011); M. H. Daouda, M. E. Rodrigues, M. J. S. Houndjo, arXiv:1109.0528 [gr-qc]; C. Q. Geng, C. C. Lee, E. N. Saridakis, Y. P. Wu, Phys. Lett. **B704**, 384 (2011) arXiv:1109.1092 [hep-th]; K. Bamba, C. Q. Geng, arXiv:1109.1694 [gr-qc]; H. Wei, arXiv:1109.6107 [gr-qc]; C. Q. Geng, C. C. Lee, E. N. Saridakis, arXiv:1110.0913 [astro-ph.CO].
- [9] B. Li, T. P. Sotiriou, J. D. Barrow, Phys. Rev. D **83**, 064035 (2011).
- [10] M. Li, R. X. Miao, Y. G. Miao, JHEP. **1107**, 108 (2011).
- [11] R. J. Yang, Europhys. Lett. **93**, 60001 (2011).
- [12] B. Li, T. P. Sotiriou, J. D. Barrow, Phys. Rev. D **83**, 104017 (2011).
- [13] J. W. Maluf, J. Math. Phys. **35**, 335 (1994).
- [14] A. Einstein, Sitz. Preuss. Akad. Wiss., 1928, p. 217; A. Einstein, Sitz. Preuss. Akad. Wiss., 1928, p. 224; A. Unzicker, T. Case, [arXiv:physics/0503046].
- [15] K. Hayashi, T. Shirafuji, Phys. Rev. D **19**, 3524 (1979); K. Hayashi, T. Shirafuji, Phys. Rev. D **24**, 3312 (1982), Addendum.
- [16] R. Aldrovandi and J. G. Pereira, An Introduction to Teleparallel Gravity, Instituto de Fisica Teorica, UNSEP, Sao Paulo (<http://www.ift.unesp.br/gcg/tele.pdf>).
- [17] R. Ferraro and F. Fiorini, Phys. Lett. **B702**, 75 (2011); R. Ferraro and F. Fiorini, arXiv:1106.6349 [gr-qc].
- [18] C. Germani, A. Kehagias, Phys. Rev. Lett. **105**, 011302 (2010); C. Germani, A. Kehagias, JCAP **05**, 019 (2010).
- [19] R. Arnowitt, S. Deser, and C. W. Misner, in Gravitation: an Introduction to Current Research, edited by L. Witten Wiley, New York, 1962.
- [20] J. W. Maluf, J. Math. Phys. **36**, 4242 (1995) .
- [21] J. Schwinger, Phys. Rev. **130**, 1253 (1963).
- [22] J. W. Maluf and J. F. da Rocha-Neto, Phys. Rev. D **64**, 084014 (2001).
- [23] T. P. Sotiriou, B. Li, J. D. Barrow, Phys. Rev. D **83**, 104030 (2011).

[24] V. Faraoni and S. Nadeau, Phys. Rev. D75, 023501 (2007).